

# Hereditary Breast and Gynecologic Cancers Panel

Pathogenic germline variants in multiple genes have been implicated in hereditary breast, ovarian, and endometrial cancers. Hereditary cancer predisposition is often characterized by an early age of cancer onset (typically before age 50) and multiple, multifocal, and/or related cancers in a single individual or in a closely related family member(s). This test includes analysis of several genes associated with hereditary breast and/or gynecologic cancer(s) that cause variable phenotypes and cancer risks, including nonbreast/nongynecologic cancers. See [Genes Tested](#) table below for more details regarding the genes and syndromes included on the Hereditary Breast and Gynecological Cancers Panel. Genes included on this panel are also included on other related tests (see [Related Tests](#) section and [Hereditary Cancer Genetic Testing – Germline Testing for Inherited Cancer Syndromes](#)).

## Disease Overview

### Associated Disorders

- *BRCA1* and *BRCA2*-associated HBOC syndrome<sup>1,2</sup>
  - Caused by a single pathogenic *BRCA1* or *BRCA2* variant
  - Individuals are at increased risk for breast, ovarian, fallopian tube, peritoneal, pancreatic, prostate, melanoma, and other cancers
  - *BRCA1* and *BRCA2* analysis is also offered through the *BRCA1* and *BRCA2*-Associated HBOC Syndrome Panel ([3001855](#))
- Lynch syndrome<sup>3</sup>
  - Caused by a single pathogenic variant in one of the mismatch repair (MMR) genes (*MLH1*, *MSH2*, *MSH6*, *PMS2*) or *EPCAM* exon 9 deletions
  - Individuals are at an increased risk for colorectal, uterine, ovarian, and other cancers
  - The Lynch syndrome genes are also offered through the Lynch Syndrome Panel, Sequencing and Deletion/Duplication ([3001605](#)). For more information, see the [Lynch Syndrome Panel, Sequencing and Deletion Test Fact Sheet](#).
- Other associated disorders on this panel include: Cowden syndrome, Li-Fraumeni syndrome (LFS), Peutz-Jeghers syndrome (PJS), and others. Please see [Genes Tested](#) table for more information.

## Genetics

### Genes

See [Genes Tested](#) table for genes included in the panel.

### Etiology

Approximately 5-10% of all breast cancers, 10-15% of ovarian cancers, and 5% of endometrial cancers are associated with a hereditary cause.<sup>4,5,6,7</sup>

## Tests to Consider

To compare directly to other hereditary cancer panels offered by ARUP Laboratories, see the [Hereditary Cancer Panel Comparison](#) table.

### [Hereditary Breast and Gynecological Cancers Panel, Sequencing and Deletion/Duplication 2012026](#)

**Method:** Massively Parallel Sequencing/Sequencing/Multiplex Ligation-dependent Probe Amplification

- Multigene panel to confirm a hereditary cause of breast and/or gynecologic cancer(s) in individuals with a complex personal or family history of breast, ovarian, or endometrial cancer
- Testing minors for adult-onset conditions is not recommended and will not be performed on minors without prior approval; for additional information, please contact an ARUP genetic counselor (800-242-2787).

### [Familial Targeted Sequencing 3005867](#)

**Method:** Massively Parallel Sequencing

- Testing for a known familial sequence variant by sequencing gene of interest. A copy of the family member's test result documenting the familial gene variant is REQUIRED.
- To determine if the variant(s) of interest are detectable by this assay, contact an ARUP genetic counselor at 800-242-2787.

See [Related Tests](#)

## Prevalence

- 1/400 individuals from general population or 1/40 Ashkenazi Jewish individuals have a *BRCA1* or *BRCA2* pathogenic variant<sup>8,9</sup>
- Lynch syndrome occurs in approximately 1/279 individuals in the general population<sup>10</sup>

## Inheritance

- Autosomal dominant
- Some genes are also associated with a predisposition to autosomal recessive childhood cancer or other syndromes.
- See [Genes Tested](#) table for additional details.

## Test Description

### Contraindications for Ordering

- Should not be ordered to detect somatic variants associated with malignancy as sensitivity for mosaic variants is low with the methodology used for germline assays
- Individuals with a hematologic malignancy and/or a previous allogeneic bone marrow transplant should not undergo molecular genetic testing on a peripheral blood specimen.
  - Testing of cultured fibroblasts is required for accurate interpretation of test results for these individuals.

### Methodology

This test is performed using the following sequence of steps:

- Selected genomic regions, primarily coding exons and exon-intron boundaries, from the targeted genes are isolated from extracted genomic DNA using a probe-based hybrid capture enrichment workflow.
- Enriched DNA is sequenced by massively parallel sequencing (MPS; also known as next generation sequencing [NGS]) followed by paired-end read alignment and variant calling using a custom bioinformatics pipeline. The pipeline includes an algorithm for detection of large (single exon-level or larger) deletions and duplications.
- Sanger sequencing is performed as necessary to fill in regions of low coverage and in certain situations, to confirm variant calls.
- Large deletion/duplication calls made using MPS are confirmed by an orthogonal exon-level microarray when sample quality and technical conditions allow.
- Long-range PCR followed by nested Sanger sequencing is performed on the following gene and exons:
  - *PMS2* (NM\_000535) 11, 12, 13, 14, 15
- Bidirectional Sanger sequencing is performed on the following genes and exons:
  - *MSH2* (NM\_000251) 5
  - *PMS2* (NM\_000535) 7
  - *PTEN* (NM\_000314) 9
- Multiplex ligation-dependent probe amplification (MLPA) is performed on the following gene to call exon-level deletions and duplications:
  - *PMS2* (NM\_000535)

### Clinical Sensitivity

Variable, dependent on phenotype/condition

- *BRCA1* and *BRCA2* sequencing and deletion/duplication testing alone detects 20-60% of hereditary breast and ovarian cancers, in general.<sup>1,4,11</sup>
- The majority of inherited endometrial cancers are thought to be caused by Lynch syndrome.

### Analytic Sensitivity

- Sanger sequencing and multiplex ligation-dependent probe amplification (MLPA) of *PMS2*: 99%
- For massively parallel sequencing:

Variant Class	Analytic Sensitivity (PPA) Estimate <sup>a</sup> (%) and 95% Credibility Region	Analytic Specificity (NPA) Estimate (%)
SNVs	>99 (96.9-99.4)	>99.9
Deletions 1-10 bp <sup>b</sup>	93.8 (84.3-98.2)	>99.9
Insertions 1-10 bp <sup>b</sup>	94.8 (86.8-98.5)	>99.9
Exon-level <sup>c</sup> deletions	97.8 (90.3-99.8) [2 exons or larger] 62.5 (38.3-82.6) [single exon]	>99.9
Exon-level <sup>c</sup> duplications	83.3 (56.4-96.4) [3 exons or larger]	>99.9
Exon-level deletions/duplications (MLPA)	>99	>99.9

<sup>a</sup>PPA values are derived from larger methods-based MPS and/or Sanger validations. These values do not apply to testing performed by MLPA unless otherwise indicated.

<sup>b</sup>Variants greater than 10 bp may be detected, but the analytic sensitivity may be reduced.

<sup>c</sup>In most cases, a single exon deletion or duplication is less than 450 bp and 3 exons span a genomic region larger than 700 bp.

bp, base pairs; NPA, negative percent agreement; PPA, positive percent agreement; SNVs, single nucleotide variants

## Limitations

- A negative result does not exclude a heritable form of cancer.
- Diagnostic errors can occur due to rare sequence variations.
- Interpretation of this test result may be impacted if this individual has had an allogeneic stem cell transplantation.
- The following will not be evaluated:
  - Variants outside the coding regions and intron-exon boundaries of the targeted genes
  - Regulatory region variants and deep intronic variants
  - Breakpoints of large deletions/duplications
  - Deletions/duplications in *NF1*, *RECQL*
  - Sequence variants in *EPCAM*
  - Noncoding transcripts
  - The following exons are not sequenced due to technical limitations of the assay:
    - *BRCA1* (NM\_007300) 13
    - *CHEK2* (NM\_001005735) 3; (NM\_001349956) 4
    - *RECQL* (NM\_002907) 14, 15; (NM\_032941) 15, 16
- The following may not be detected:
  - Deletions/duplications/insertions of any size by MPS
  - Large duplications less than 3 exons in size
  - Noncoding transcripts
  - Single exon deletions/duplications may not be detected based on the breakpoints of the rearrangement
  - Some variants due to technical limitations in the presence of pseudogenes and/or repetitive/homologous regions
  - Low-level somatic variants
  - Deletions/duplications in the following exons:

Gene	Exon(s)
<i>BRCA1</i>	(NM_007294, NM_007299, NM_007300) 2; (NM_007298) 1
<i>CDH1</i>	(NM_001317185) 10

Gene	Exon(s)
<i>CHEK2</i>	(NM_007194) 11-15; (NM_001005735) 3,12-16; (NM_001257387) 12-16; (NM_001349956) 4,10-14; (NM_145862) 10-14
<i>PTEN</i>	(NM_000314, NM_001304718) 9; (NM_001304717) 1,10
<i>RECQL</i>	(NM_002907) 14-15; (NM_032941) 15-16

## Genes Tested

To compare directly to other hereditary cancer panels offered by ARUP Laboratories, see the [Hereditary Cancer Panel Comparison table](#).

Gene	MIM Number	Disorder	Inheritance
<i>ATM</i>	607585	Breast, colorectal, <sup>a</sup> ovarian, pancreas, prostate	AD
		Ataxia-telangiectasia	AR
<i>BARD1</i>	601593	Breast <sup>a</sup>	AD
<i>BRCA1</i>	113705	HBOC syndrome Breast, fallopian tube, melanoma, ovarian, pancreatic, peritoneal, prostate	AD
		Fanconi anemia, complementation group S	AR
<i>BRCA2</i>	600185	HBOC syndrome Breast, fallopian tube, melanoma, ovarian, pancreatic, peritoneal, prostate	AD
		Fanconi anemia, complementation group D1	AR
<i>BRIP1</i>	605882	Breast, <sup>a</sup> ovarian	AD
		Fanconi anemia, complementation group J	AR
<i>CDH1</i>	192090	HDGC Diffuse gastric, lobular breast	AD
<i>CHEK2</i>	604373	Breast, colorectal, prostate, thyroid <sup>a</sup>	AD
<i>DICER1</i>	606241	<i>DICER1</i> -related disorders CNS, cystic nephroma, ovarian sex cord-stromal tumors, pleuropulmonary blastoma, thyroid	AD

<sup>a</sup>Association is suggested but not well-established at this time.

AD, autosomal dominant; AR, autosomal recessive; CMMRD, constitutional mismatch repair deficiency; CNS, central nervous system; GIST, gastrointestinal stromal tumor; HBOC, hereditary breast and ovarian cancer; HDGC, hereditary diffuse gastric cancer; HNPCC, hereditary nonpolyposis colorectal cancer; LFS, Li-Fraumeni syndrome; NBS, Nijmegen breakage syndrome; NF1, neurofibromatosis type 1; PJS, Peutz-Jeghers syndrome; RTPS, rhabdoid tumor predisposition syndrome; SCCOHT, small-cell carcinoma of the ovary—hypercalcemic type

Gene	MIM Number	Disorder	Inheritance
<b>EPCAM</b> (Exon 9 deletions/duplications only)	185535	Lynch syndrome/HNPCC Brain, colorectal, endometrial, ovarian, pancreas, prostate, renal pelvis and/or ureter, stomach, and others	AD
<b>MLH1</b>	120436	Lynch syndrome/HNPCC Brain, colorectal, endometrial, ovarian, pancreas, prostate, renal pelvis and/or ureter, stomach, and others	AD
		CMMRD	AR
<b>MSH2</b>	609309	Lynch syndrome/HNPCC Brain, colorectal, endometrial, ovarian, pancreas, prostate, renal pelvis and/or ureter, stomach, and others	AD
		CMMRD	AR
<b>MSH6</b>	600678	Lynch syndrome/HNPCC Brain, colorectal, endometrial, ovarian, pancreas, prostate, renal pelvis and/or ureter, stomach, and others	AD
		CMMRD	AR
<b>NBN</b>	602667	Breast, <sup>a</sup> ovarian, <sup>a</sup> prostate <sup>a</sup>	AD
		NBS	AR
<b>NF1</b>	613113	NF1 Breast, GIST, gliomas, leukemia, malignant peripheral nerve sheath tumors, neurofibromas, pheochromocytoma	AD
<b>PALB2</b>	610355	Breast, ovarian, pancreas, prostate	AD
		Fanconi anemia, complementation group N	AR
<b>PMS2</b>	600259	Lynch syndrome/HNPCC Brain, colorectal, endometrial, ovarian, pancreas, prostate, renal pelvis and/or ureter, stomach, and others	AD
		CMMRD	AR
<b>PTEN</b>	601728	Cowden syndrome/ <i>PTEN</i> hamartoma tumor syndrome Breast, colorectal, endometrial, Lhermitte-Duclos disease (cerebellar dysplastic gangliocytoma), melanoma, <sup>a</sup> renal cell carcinoma, thyroid, and others	AD
<b>RAD51C</b>	602774	Breast, ovarian	AD

<sup>a</sup>Association is suggested but not well-established at this time.

AD, autosomal dominant; AR, autosomal recessive; CMMRD, constitutional mismatch repair deficiency; CNS, central nervous system; GIST, gastrointestinal stromal tumor; HBOC, hereditary breast and ovarian cancer; HDGC, hereditary diffuse gastric cancer; HNPCC, hereditary nonpolyposis colorectal cancer; LFS, Li-Fraumeni syndrome; NBS, Nijmegen breakage syndrome; NF1, neurofibromatosis type 1; PJS, Peutz-Jeghers syndrome; RTPS, rhabdoid tumor predisposition syndrome; SCCOHT, small-cell carcinoma of the ovary–hypercalcemic type

Gene	MIM Number	Disorder	Inheritance
		Fanconi anemia, complementation group O	AR
<b>RAD51D</b>	602954	Breast, ovarian, prostate	AD
<b>RECQL</b>	600537	Breast <sup>a</sup>	AD
<b>SMARCA4</b>	603254	Coffin-Siris syndrome, RTPS Rhabdoid tumors located in CNS, kidney, ovary (SCCOHT), and others	AD
<b>STK11</b>	602216	PJS Breast, cervix, colorectal, endometrial, lung, ovarian (sex cord with annular tubules), pancreas, Peutz-Jeghers-type hamartomatous polyps, small intestine, stomach, testes	AD
<b>TP53</b>	191170	LFS Adrenocortical carcinoma, breast, choroid plexus carcinoma, CNS, colorectal, melanoma, <sup>a</sup> osteosarcoma, pancreas, prostate, renal, rhabdomyosarcoma, soft tissue sarcoma, stomach, thyroid, and others	AD

<sup>a</sup>Association is suggested but not well-established at this time.

AD, autosomal dominant; AR, autosomal recessive; CMMRD, constitutional mismatch repair deficiency; CNS, central nervous system; GIST, gastrointestinal stromal tumor; HBOC, hereditary breast and ovarian cancer; HDGC, hereditary diffuse gastric cancer; HNPCC, hereditary nonpolyposis colorectal cancer; LFS, Li-Fraumeni syndrome; NBS, Nijmegen breakage syndrome; NF1, neurofibromatosis type 1; PJS, Peutz-Jeghers syndrome; RTPS, rhabdoid tumor predisposition syndrome; SCCOHT, small-cell carcinoma of the ovary—hypercalcemic type

## References

- Petrucelli N, Daly MB, Pal T. [BRCA1- and BRCA2-associated hereditary breast and ovarian cancer](#). In: Adam MP, Ardinger HH, Pagon RA, et al, editors. GeneReviews, University of Washington; 1993-2022. [Last update: Feb 2022; Accessed: Feb 2022]
- National Comprehensive Cancer Network. [NCCN Clinical Practice Guidelines in Oncology: genetic/familial high-risk assessment—breast, ovarian, and pancreatic](#). Version 2.2021. [Updated: Nov 2020; Accessed: Aug 2021]
- National Comprehensive Cancer Network. [NCCN Clinical Practice Guidelines in Oncology: genetic/familial high-risk assessment—colorectal](#). Version 1.2020. [Updated: Jul 2020; Accessed: Feb 2021]
- Pruthi S, Gostout BS, Lindor NM. [Identification and management of women with BRCA mutations or hereditary predisposition for breast and ovarian cancer](#). *Mayo Clin Proc*. 2010;85(12):1111-1120.
- Couch FJ, Shimelis H, Hu C, et al. [Associations between cancer predisposition testing panel genes and breast cancer](#). *JAMA Oncol*. 2017;3(9):1190-1196.
- Risch HA, McLaughlin JR, Cole DE, et al. [Prevalence and penetrance of germline BRCA1 and BRCA2 mutations in a population series of 649 women with ovarian cancer](#). *Am J Hum Genet*. 2001;68(3):700-710.
- National Comprehensive Cancer Network. [NCCN Clinical Practice Guidelines in Oncology: Uterine neoplasms](#). Version 1.2022. [Updated: Nov 2021; Accessed: May 2022]
- Whittemore AS, Gong G, John EM, et al. [Prevalence of BRCA1 mutation carriers among U.S. non-Hispanic Whites](#). *Cancer Epidemiol Biomarkers Prev*. 2004;13(12):2078-2083.
- King MC, Marks JH, Mandell JB, et al. [Breast and ovarian cancer risks due to inherited mutations in BRCA1 and BRCA2](#). *Science*. 2003;302(5645):643-646.
- Win AK, Jenkins MA, Dowty JG, et al. [Prevalence and penetrance of major genes and polygenes for colorectal cancer](#). *Cancer Epidemiol Biomarkers Prev*. 2017;26(3):404-412.
- Meindl A, Ditsch N, Kast K, et al. [Hereditary breast and ovarian cancer: new genes, new treatments, new concepts](#). *Dtsch Arztebl Int*. 2011;108(19):323-330.

## Additional Resources

Chen S, Wang W, Lee S, et al. [Prediction of germline mutations and cancer risk in the Lynch syndrome](#), JAMA. 2006;296(12):1479-1487.

Doros L, Schultz KA, Stewart DR, et al. [DICER1-related disorders](#). In: Adam MP, Ardinger HH, Pagon RA, et al, eds. GeneReviews, University of Washington; 1993-2021. [Last update: Apr 2020; Accessed: Jun 2020]

Ford D, Easton DF, Stratton M, et al. [Genetic heterogeneity and penetrance analysis of the BRCA1 and BRCA2 genes in breast cancer families. The Breast Cancer Linkage Consortium](#). *Am J Hum Genet*. 1998;62(3):676-689.

Friedman JM. [Neurofibromatosis 1](#). In: Adam MP, Ardinger HH, Pagon RA, et al., eds. GeneReviews, University of Washington; 1993-2021. [Updated: Apr 2022; Accessed: May 2022]

Laloo F, Evans DG. [Familial breast cancer](#). *Clin Genet*. 2012;82(2):105-114.

Nemes K, Bens S, Bourdeaut F, et al. [Rhabdoid tumor predisposition syndrome](#). In: Adam MP, Ardinger HH, Pagon RA, et al, editors. GeneReviews, University of Washington; 1993-2020. [Initial Posting: Dec 2017; Accessed: Feb 2020]

Walsh T, Casadei S, Lee MK, et al. [Mutations in 12 genes for inherited ovarian, fallopian tube, and peritoneal carcinoma identified by massively parallel sequencing](#). *Proc Natl Acad Sci U S A*. 2011;108(44):18032-18037.

## Related Information

[Breast Cancer Biomarkers](#)  
[Hereditary Cancer Germline Genetic Testing](#)  
[Ovarian Cancer Biomarkers](#)

## Related Tests

[BRCA1 and BRCA2-Associated HBOC Syndrome Panel, Sequencing and Deletion/Duplication 3001855](#)

**Method:** Massively Parallel Sequencing

[Lynch Syndrome Panel, Sequencing and Deletion/Duplication 3001605](#)

**Method:** Massively Parallel Sequencing/Sequencing/Multiplex Ligation-dependent Probe Amplification

[Hereditary Cancer Panel, Sequencing and Deletion/Duplication 2012032](#)

**Method:** Massively Parallel Sequencing/Sequencing/Multiplex Ligation-dependent Probe Amplification

[Neurofibromatosis Type 1 and Legius Syndrome Panel, Sequencing and Deletion/Duplication 3003927](#)

**Method:** Massively Parallel Sequencing

[Hereditary Breast Cancer Guidelines-Based Panel, Sequencing and Deletion/Duplication 3005654](#)

**Method:** Massively Parallel Sequencing/Sequencing

[Hereditary Breast Cancer High-Risk Panel, Sequencing and Deletion/Duplication 3005632](#)

**Method:** Massively Parallel Sequencing/Sequencing

ARUP Laboratories is a nonprofit enterprise of the University of Utah and its Department of Pathology. 500 Chipeta Way, Salt Lake City, UT 84108  
(800) 522-2787 | (801) 583-2787 | aruplab.com | arupconsult.com  
Content Review May 2022 | Last Update November 2022