

# Autoimmune Neurologic Disease Panel With Reflex, Serum and CSF

Last Literature Review: May 2023 | Last Update: September 2024

Autoimmune neurologic diseases represent a broad category of conditions characterized by immune dysregulation. Antibodies associated with these conditions may be present in the serum or cerebrospinal fluid (CSF) and can serve as useful markers of disease. Testing for these antineural antibodies should be considered when patients experience subacute onset of new, unexplained neurologic symptoms.

## Disease Overview

Autoimmune neurologic diseases may localize to the peripheral nervous system or central nervous system (CNS). They can manifest with diverse symptoms including (but not limited to) brainstem or cerebellar syndromes, dysautonomia, encephalopathy, epilepsy, movement disorders, myelopathy, psychiatric changes, or rapidly progressive dementia. Antineural antibodies serve as useful markers of these diseases, and their detection may help establish a diagnosis, support treatment decisions, aid prognostication, serve as a prerequisite for enrollment in clinical trials, and guide the search for an associated malignancy.

For more information about laboratory testing for autoimmune neurologic diseases, including detailed information about panel test selection, refer to the ARUP Consult [Autoimmune Neurologic Diseases - Antineural Antibody Testing](#) topic.

## Test Description

ARUP's serum or CSF Autoimmune Neurologic Disease With Reflex panels can be used for the evaluation of patients with subacute onset of neurologic symptoms with possible autoimmune etiology. Testing for the presence of antineural antibodies in both serum and CSF is recommended in most situations.<sup>1</sup>

These panels cover a broad range of autoimmune neurologic phenotypes; for greater diagnostic yield and improved turnaround time, consider choosing a phenotype-specific panel (see table below) rather than a broad panel.

ARUP Phenotype-Specific Panels to Consider for Autoimmune Neurologic Disease		
ARUP Panel	Test Code	
	Serum	CSF
Autoimmune Encephalopathy/Dementia Panel	<a href="#">3006201</a>	<a href="#">3006202</a>
Autoimmune Epilepsy Panel	<a href="#">3006204</a>	<a href="#">3006205</a>
Autoimmune Movement Disorder Panel	<a href="#">3006206</a>	<a href="#">3006207</a>
Autoimmune Myelopathy Panel	<a href="#">3006208</a>	<a href="#">3006209</a>
Autoimmune Dysautonomia Panel	<a href="#">3006203</a>	—
Autoimmune Pediatric CNS Disorders	<a href="#">3006210</a>	<a href="#">3006211</a>

Regardless of the panel chosen, order only one panel for serum and/or one panel for CSF; many antineural antibodies are redundant between these panels, and choosing based on the predominant phenotype will provide the most meaningful results. To compare these panels and the antibodies included, refer to the ARUP [Antineural Antibody Testing for Autoimmune Neurologic Disease](#) page.

Testing for individual autoantibodies is also available separately and can be used for long-term monitoring.

### Featured ARUP Testing

#### [Autoimmune Neurologic Disease Panel with Reflex, Serum 3006051](#)

**Method:** Semi-Quantitative Cell-Based Indirect Fluorescent Antibody/Qualitative Immunoblot/Quantitative Radioimmunoassay (RIA)/Semi-Quantitative Enzyme-Linked Immunosorbent Assay (ELISA)

#### [Autoimmune Neurologic Disease Panel With Reflex, CSF 3006052](#)

**Method:** Semi-Quantitative Cell-Based Indirect Fluorescent Antibody/Qualitative Immunoblot/Quantitative Radioimmunoassay (RIA)/Semi-Quantitative Enzyme-Linked Immunosorbent Assay (ELISA)

## Antibodies Tested and Methodology

Autoimmune Neurologic Disease Panel With Reflex, Serum (3006051) and CSF (3006052): Antibodies Tested and Methodology			
Autoantibody Marker	Method	Individual Autoantibody or Focused Panel Test Code	
		Serum	CSF
AChR binding Ab, IgG <sup>a</sup>	RIA	0080009	—
AMPA Ab, IgG	CBA-IFA	3001260	3001257
Amphiphysin Ab, IgG	IB	2008893	3004510
ANNA-1 (Hu)	IFA, reflex IB, reflex titer	2007961	2010841
ANNA-2 (Ri)	IFA, reflex IB, reflex titer	2007961	2010841
AQP4 Ab, IgG	CBA-IFA, reflex titer	2013320	2011699 <sup>b</sup>
CASPR2 Ab, IgG	CBA-IFA, reflex titer	2009452	3001986
CV2 (CRMP-5) Ab, IgG	CBA-IFA, reflex titer	3016999	3017001
DPPX Ab, IgG	CBA-IFA, reflex titer	3004359	3004512
GABA-AR Ab, IgG	CBA-IFA, reflex titer	3006008	3006003
GABA-BR Ab, IgG	CBA-IFA, reflex titer	3001270	3001267
GAD65 Ab	ELISA	2001771	3002788
Ganglionic AChR Ab, IgG	RIA	3003020	—
IgLON5 Ab, IgG	CBA-IFA, reflex titer	3006018	3006013
ITPR1 Ab, IgG	CBA-IFA, reflex titer	3006031	3006023
LG11 Ab, IgG	CBA-IFA, reflex titer	2009456	3001992
mGluR1 Ab, IgG	CBA-IFA, reflex titer	3006044	3006039
MOG Ab, IgG	CBA-IFA, reflex titer	3001277	—
NMDAR Ab, IgG	CBA-IFA, reflex to titer	2004221	2005164
PCCA-1 (Yo)	IFA, reflex IB, reflex titer	2007961	2010841
PCCA-Tr/DNER	IFA, reflex IB, reflex titer	2007961	2010841
P/Q-type VGCC Ab, IgG	RIA	0092628	—
SOX1 (AGNA) Ab, IgG	IB	3002885	3002886
VGKC Ab, IgG	RIA	2004890	3001387


<sup>a</sup>Performed via reflex only, depending on the results of other autoantibody tests; refer to Reflex Patterns flowchart.

<sup>b</sup>Testing for AQP4 Ab is not included in the Autoimmune Neurologic Disease Panel With Reflex, CSF panel.

Ab, antibody; AChR, acetylcholine receptor; AGNA, antiglial nuclear antibody; AMPAR, alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor; ANNA, antineuronal nuclear antibody; AQP4, aquaporin-4; CASPR2, contactin-associated protein 2; CBA, cell-binding assay/cell-based assay; CRMP-5, collapsin response-mediator protein 5; DNER, Delta/notch-like epidermal growth factor-related receptor; DPPX, dipeptidyl-aminopeptidase-like protein 6; GABA-AR, gamma-aminobutyric acid receptor, type A; GABA-BR, gamma aminobutyric acid receptor, type B; GAD65, glutamic acid decarboxylase antibody 65-kd isoform; IFA, indirect immunofluorescence assay; Ig, immunoglobulin; IgLON5; IgLON family member 5; ITPR1, inositol 1,4,5-triphosphate receptor type 1; LG11, leucine-rich, glioma-inactivated protein 1; mGluR1, metabotropic glutamate receptor 1; MOG, myelin oligodendrocyte glycoprotein; NMDAR, N-methyl-D-aspartate receptor antibody; PCCA, Purkinje cell cytoplasmic antibody; RIA, radioimmunoassay; SOX1, SRY-box transcription factor 1; VGCC, voltage gated calcium channel; VGKC, voltage-gated potassium channel

## Reflex Patterns

Autoimmune Neurologic Disease Panel With Reflex, Serum (3006051) and CSF (3006052)



## Limitations

These tests do not include all known antineural antibodies:

- Some antibodies are extremely rare or are of uncertain clinical significance.
- As testing for newly described antibodies becomes available and their clinical relevance is established, these panels will evolve to reflect these discoveries.

## Test Interpretation

### Results

Results must be interpreted in the clinical context of the individual patient; test results (positive or negative) should not supersede clinical judgement.

Autoimmune Neurologic Disease Panel With Reflex, Serum (3006051) and CSF (3006052): Results Interpretation	
Result	Interpretation
Positive for ≥1 autoantibodies	Autoantibody(ies) detected May support a diagnosis of autoimmune neurologic disease Consider a focused search for malignancy based on antibody-tumor associations
Negative	No autoantibodies detected A diagnosis of autoimmune neurologic disease is not excluded

## References

1. Budhram A, Dubey D, Sechi E, et al. [Neural antibody testing in patients with suspected autoimmune encephalitis](#). *Clin Chem*. 2020;66(12):1496-1509.

ARUP Laboratories is a nonprofit enterprise of the University of Utah and its Department of Pathology. 500 Chipeta Way, Salt Lake City, UT 84108  
(800) 522-2787 | (801) 583-2787 | aruplab.com | arupconsult.com

© 2024 ARUP Laboratories. All Rights Reserved.

Client Services - (800) 522-2787